

A conjectured form of the Goldberg-Sachs theorem¹

Andrzej Trautman

Instytut Fizyki Teoretycznej, Uniwersytet Warszawski
Hoża 69, 00681 Warszawa, Poland
e-mail: amt@fuw.edu.pl

The rather well-known Goldberg-Sachs theorem [?] is one of the most beautiful results in the mathematics of general relativity theory. It played a major role in the work on algebraically special solutions of Einstein's equations [?].

For the purposes of this Letter, it is convenient to formulate it as follows. Let \mathfrak{M} be a set of Lorentzian, not conformally flat, manifolds (M, g) of dimension 4. For $(M, g) \in \mathfrak{M}$, let $K \subset TM$ be a null line bundle; its sections are null vector fields. Following the notation and terminology of [?], consider the following two properties of K :

(GSR) K is geodetic and shear-free;

(PND) K is a bundle of *repeated* principal null directions of the Weyl tensor C .

A Goldberg-Sachs theorem $\text{GST}(\mathfrak{M})$ is a statement of the form: *if $(M, g) \in \mathfrak{M}$, then the conditions (GSR) and (PND) are equivalent.* Goldberg and Sachs proved the theorem for $\mathfrak{M} =$ the set of Einstein spaces, i.e. solutions of $R_{\mu\nu} = \frac{1}{4}g_{\mu\nu}R$. Shortly afterwards, Kundt and Thompson [?] and Robinson and Schild [?] pointed out that both conditions (GSR) and (PND) are conformally invariant, but the property of being an Einstein space is not. They proved a generalized Goldberg-Sachs theorem that, in a refined form, is given in §7.3 of [?] and in §7.5 of [?]. This generalized theorem involves only conformal notions, but requires a separate formulation for each degree of degeneracy of the Weyl tensor.

A conformally invariant set of space-times is

$$\mathfrak{M}_c = \{(M, g) \text{ is conformal to an Einstein space}\}$$

and $\text{GST}(\mathfrak{M}_c)$ is true as a consequence of the classical Goldberg-Sachs theorem. It is not easy to find a description of \mathfrak{M}_c by means of tensorial or spinorial equations; see [?] for an account of the early work by Brinkman. To appreciate the difficulty of the subject, recall that Schouten (p. 314 in [?]) attributes to Brinkman the following statement: In dimension 4, if two manifolds are Ricci flat and conformal to each other, but not to a flat space, then they are isometric. Ehlers and Kundt (p. 99 in [?]) give a counterexample to this: there are *pp* waves that are conformal, but not isometric, to each other.

It is known that \mathfrak{M}_c is contained in the set \mathfrak{M}_b of spaces satisfying the conformally invariant *Bach equation* $B = 0$. According to the arguments due to Geroch and Horowitz, presented in [?], there are space-times satisfying the Bach equation that are not conformal to an Einstein space.

¹During the *Workshop on spinors and twistors* (28 June - 3 July 1999) at the Erwin Schrödinger Institute in Vienna I obtained valuable advice on the subject of this note from M. Dunajski, C. N. Kozameh, L. J. Mason, P. Nurowski, and K. P. Tod. When preparing the Letter, I have been supported in part by the Polish Committee for Scientific Research (KBN) under grant no. 2 P03B 060 17.

Published in *Twistor Newsletter* **45** 50-51 (2000).

Kozameh, Newman, and Tod [?] have found a set of two equations defining a class of spaces conformal to Einstein spaces; one of them is $B = 0$, but the other one excludes some of the spaces with a degenerate C . There is an improvement of [?] by Baston and Mason [?], but their equations still do not characterize all of \mathfrak{M}_c .

Problems

I consider the following problems to be ordered according to increasing difficulty.

- (i) Find a counterexample to $\text{GST}(\mathfrak{M}_b)$.
- (ii) If you fail in (i), then prove $\text{GST}(\mathfrak{M}_b)$.
- (iii) If you succeed in (i), then find a set of conformally invariant tensor or spinor equations defining \mathfrak{M} , without reference to the degeneracy of C , such that $\mathfrak{M}_c \subset \mathfrak{M}$ and $\text{GST}(\mathfrak{M})$ is true.

Note that if $\mathfrak{M}_c \subsetneq \mathfrak{M}$, then $\text{GST}(\mathfrak{M})$ is stronger than the classical theorem.

References

- [1] R. J. Baston and L. J. Mason, Conformal gravity, the Einstein equations and spaces of complex null geodesics *Class. Quantum Grav.* **4** (1987) 815–26.
- [2] J. Ehlers and W. Kundt, Ch. 2 in *Gravitation: an introduction to current research*, ed. by L. Witten, Wiley, New York 1962.
- [3] J. N. Goldberg and R. K. Sachs, A theorem on Petrov types *Acta Phys. Polon.* **22** Suppl. (1962) 13–23.
- [4] C. N. Kozameh, E. T. Newman and K. P. Tod, Conformal Einstein spaces *Gen. Rel. Grav.* **17** (1985) 343–52.
- [5] D. Kramer, H. Stephani, M. MacCallum and E. Herlt, *Exact solutions of Einstein's field equations* VEB Deutscher Verlag der Wissenschaften, Berlin 1980.
- [6] W. Kundt and A. Thompson, Le tenseur de Weyl et une congruence associée de géodésiques isotropes sans distorsion *C. R. Acad. Sci. Paris* **254** (1962) 4257–9.
- [7] R. Penrose and W. Rindler, *Spinors and space-time* vol. 2, C. U. P., Cambridge 1986.
- [8] I. Robinson and A. Schild, Generalization of a theorem by Goldberg and Sachs *J. Math. Phys.* **4** (1963) 484–9.
- [9] J. A. Schouten, *Ricci-Calculus* 2nd ed., Springer, Berlin 1954.